

Double-Spin Asymmetry A_{LT} for Polarized Drell-Yan Process in $p\bar{p}$ Collisions: Wandzura-Wilczek Contribution

Yuji Koike*, Kazuhiro Tanaka[†] and Shinsuke Yoshida*

*Department of Physics, Niigata University, Ikarashi, Niigata 950-2181, Japan

[†]Department of Physics, Juntendo University, Inba, Chiba 270-1695, Japan

Abstract. The longitudinal-transverse spin asymmetry A_{LT} in the polarized Drell-Yan process depends on twist-3 quark distributions of nucleon. In addition to the contributions associated with the twist-3 operators, these distributions contain the “Wandzura-Wilczek (WW)” part, which is determined by a certain integral of twist-2 distributions. The recently obtained empirical information on the transversity allows a realistic estimate of the WW contribution to A_{LT} for the polarized $p\bar{p}$ collisions. Our results indicate that rather large A_{LT} ($\sim 10\%$) can be observed in the proposed spin experiments at GSI, with novel pattern as a function of dilepton mass compared with A_{TT} and A_{LL} .

Keywords: Drell-Yan process, Antiprotons, Longitudinal-transverse spin asymmetry, Twist-3

PACS: 12.38.-t, 13.85.Qk, 13.88.+e

The proposed polarization experiments with antiprotons at GSI provide promising opportunities to probe chiral-odd spin-dependent parton distributions h_1 , h_L [1]. The double transverse-spin asymmetry for the Drell-Yan (DY) process in collisions of transversely polarized protons and antiprotons, $p^\uparrow \bar{p}^\uparrow \rightarrow l^+ l^- X$, is given as ($d\omega \equiv dQ^2 dx_F d\Omega$)

$$A_{TT} = \frac{d\sigma^{\uparrow\uparrow}/d\omega - d\sigma^{\uparrow\downarrow}/d\omega}{d\sigma^{\uparrow\uparrow}/d\omega + d\sigma^{\uparrow\downarrow}/d\omega} = \hat{a}_{TT}(\theta, \phi) \frac{\sum_a e_a^2 h_1^a(x_1, Q^2) h_1^a(x_2, Q^2)}{\sum_a e_a^2 f_1^a(x_1, Q^2) f_1^a(x_2, Q^2)}, \quad (1)$$

at the leading order (LO) QCD, for the dilepton production with the invariant mass Q and the direction given by the angle $\Omega = (\theta, \phi)$. h_1^a and f_1^a denote the transversity and unpolarized quark-distributions inside proton, and the summation is over all quark and anti-quark flavors with e_a the corresponding electric charge. The scaling variables $x_{1,2}$ represent the momentum fractions associated with the partons annihilating via the DY mechanism, such that $Q^2 = (x_1 P_1 + x_2 P_2)^2 = x_1 x_2 s$ and $x_F = x_1 - x_2$, where $s = (P_1 + P_2)^2$ is the CM energy squared of $p^\uparrow \bar{p}^\uparrow$. $\hat{a}_{TT}(\theta, \phi)$ represents the asymmetry in the parton level. In particular, moderate energies at GSI, $30 \lesssim s \lesssim 200 \text{ GeV}^2$, allow us to measure (1) for $0.2 \lesssim Q/\sqrt{s} \lesssim 0.7$, and probe $h_1(x_{1,2}, Q^2)$ in the “valence region”. It turns out [2] that A_{TT} at GSI is quite stable when including the QCD (resummation and fixed-order) corrections, and that the recent empirical information on h_1 [3] leads to large value of (1) at GSI. We note that the double longitudinal-spin asymmetry A_{LL} for $p^\rightarrow \bar{p}^\rightarrow \rightarrow l^+ l^- X$ is given by (1) with the replacement $h_1 \rightarrow g_1$, $\hat{a}_{TT} \rightarrow \hat{a}_{LL}$, where g_1 is the chiral-even, helicity quark-distribution and $\hat{a}_{LL}$ is the corresponding partonic asymmetry.

It should not be overlooked that the double-spin longitudinal-transverse asymmetry $A_{LT} = [d\sigma^{\rightarrow\uparrow}/d\omega - d\sigma^{\rightarrow\downarrow}/d\omega]/[d\sigma^{\rightarrow\uparrow}/d\omega + d\sigma^{\rightarrow\downarrow}/d\omega]$ is also readily accessible in the DY experiments at GSI: A_{LT} plays a distinguished role in spin physics, allowing us

to access the twist-3 spin-dependent parton distributions as leading effects (see [1]),

$$A_{LT} = \frac{M}{Q} \hat{a}_{LT}(\theta, \phi) \frac{\sum_a e_a^2 [g_1^a(x_1, Q^2) x_2 g_T^a(x_2, Q^2) + x_1 h_L^a(x_1, Q^2) h_1^a(x_2, Q^2)]}{\sum_a e_a^2 f_1^a(x_1, Q^2) f_1^a(x_2, Q^2)}, \quad (2)$$

with the proton mass M , although suppressed by M/Q compared to A_{TT} , A_{LL} that receive contribution only from twist-2 distributions. g_T^a and h_L^a denote the twist-3 distributions associated with the transversely and longitudinally polarized proton, respectively; the chiral-even distribution g_T^a is also accessible by the longitudinal-transverse asymmetry associated with the structure function g_2 in the polarized DIS [4], but the chiral-odd h_L^a is not accessible by inclusive DIS. We assess the potential of A_{LT} at GSI experiments [1], extending the previous study on A_{LT} in pp -collision at RHIC energy [5].

The distributions h_L and g_T contain the piece expressed by twist-2 matrix element as

$$h_L^a(x, Q^2) = 2x \int_x^1 dy \frac{h_1^a(y, Q^2)}{y^2} + \dots, \quad g_T^a(x, Q^2) = \int_x^1 dy \frac{g_1^a(y, Q^2)}{y} + \dots, \quad (3)$$

where the ellipses stand for “genuine twist-3” contributions given as matrix element of the twist-3 operators; those twist-3 operators can be reexpressed as quark-gluon-quark three-body correlation operators on the lightcone, using the QCD equations of motion (see [6, 7]). We call the twist-2 component, shown explicitly in (3), the “Wandzura-Wilczek (WW)” part. Because the operators with different “geometric twist” do not mix with each other under QCD evolution with Q^2 , both x - and Q^2 -dependences of the WW part are determined solely by those of the corresponding twist-2 distributions.

We work at LO QCD with (2), which provides sufficient accuracy for our first estimate of A_{LT} at GSI. We may anticipate that QCD corrections to A_{LT} at GSI kinematics could be small, similarly to the case for A_{TT} mentioned above. We also employ the “WW approximation” to A_{LT} , i.e., evaluate (2) using (3) with the genuine twist-3 contributions omitted: the WW approximation for g_T of (3) is supported by the data on the structure function g_2 from the polarized DIS experiments [4], as well as from lattice QCD simulation [8]. Estimates from nucleon models, combined with the QCD evolution which is different between the relevant twist-2 [9] and twist-3 operators [6, 7] such that the latter is more strongly suppressed for high Q^2 than the former, also suggest that the WW part of (3) dominates h_L and g_T for $Q^2 \gg 1 \text{ GeV}^2$ [5, 10].

We use the LO GRV98 [11] and GRSV2000 (“standard scenario”) [12] distributions for the unpolarized and longitudinally-polarized quark distributions f_1 and g_1 , respectively. For the LO transversity distribution h_1 , we are guided by the recent information from the LO global fit [3]: we find that a useful estimate can be obtained by assuming the relation, $h_1^a(x, \mu^2) = g_1^a(x, \mu^2)$, at a low scale μ ($\mu^2 = 0.26 \text{ GeV}^2$ using the GRSV2000 g_1); its QCD evolution from μ^2 to Q^2 is controlled by the LO DGLAP kernel [9] for h_1 . Comparing the resulting transversity [1] with h_1 from the LO global fit [3], our h_1 lies slightly outside the error band of the fit, for the valence region $0.2 \lesssim x \lesssim 0.7$ relevant for the GSI kinematics. Thus our h_1 will provide a realistic estimate of the upper bound of the relevant asymmetries using (1)-(3), implied by the present empirical uncertainty in transversity. Similar choice for h_1 has been adopted to calculate QCD corrections for A_{TT} at GSI [2]. In the valence region relevant for the GSI kinematics, the dominance




FIGURE 1. (a) $\tilde{A}_{LT}$ and (b) $\tilde{A}_{LL}$ and $\tilde{A}_{TT}$ as a function of x_F for $Q = 2.5$ and 4 GeV at $s = 45$ GeV².

of the u -quark contribution is observed for all twist-2 quark-distributions, $(h_1^u)^2 \gg (h_1^a)^2$ with $a = \bar{u}, d, \bar{d}, \dots$ [1, 3], and similarly for g_1 and f_1 [12, 11]. Calculating (3) using our twist-2 distributions, we find the similar u -quark dominance for h_L and g_T in the WW approximation [1]. Therefore, from (2) at GSI kinematics,

$$\tilde{A}_{LT} \equiv \frac{A_{LT}}{(M/Q)\hat{a}_{LT}(\theta, \phi)} \simeq \frac{g_1^u(x_1, Q^2)x_2 \int_{x_2}^1 dy \frac{g_1^u(y, Q^2)}{y} + 2x_1^2 \int_{x_1}^1 dy \frac{h_1^u(y, Q^2)}{y^2} h_1^u(x_2, Q^2)}{f_1^u(x_1, Q^2) f_1^u(x_2, Q^2)} + \dots, \quad (4)$$

where the ellipses denote the contributions associated with the genuine twist-3 operators. Thus the WW approximation to A_{LT} at GSI is directly related to the behavior of h_1^u .

We present the results [1] for the “reduced asymmetries”, $\tilde{A}_{LT}$, defined in (4), and $\tilde{A}_{YY} \equiv A_{YY}/\hat{a}_{YY}$ ($Y = L, T$). Figure 1(a) shows $\tilde{A}_{LT}$, along with the separated contributions from the chiral-even and -odd distributions corresponding to the first and second terms in the numerator in (2). The value of $\tilde{A}_{LT}$ is significant. Its x_F -behavior is in strong contrast to the flat and symmetric behavior of $\tilde{A}_{LL}$, $\tilde{A}_{TT}$ at the same kinematics, shown in Fig. 1(b). $\tilde{A}_{LT}$ is smaller than $\tilde{A}_{TT}$, $\tilde{A}_{LL}$ by the additional $x_{1,2}$ factor in (2) compared with (1), and also by the behavior of h_L and g_T where the integral for the WW part, with the factors $1/y^2$, $1/y$ in (3), shifts the peak of the distributions to lower x with the suppressed peak-height, compared with the corresponding twist-2 distributions [1]. Indeed, for pp collisions where the sea-quark region is probed, these effects, in particular the additional $x_{1,2}$ factor, lead to $\tilde{A}_{LT}$ much smaller than the corresponding $\tilde{A}_{LL}$, $\tilde{A}_{TT}$ [5].

Figure 2 shows the relevant asymmetries with $x_1 = x_2 = Q/\sqrt{s}$ as a function of Q . The Q dependence of $\tilde{A}_{TT}$ as well as $\tilde{A}_{LL}$ directly reflects the x dependence of the corresponding distributions, because the u -quark dominance in (1) implies that $\tilde{A}_{TT}$ is controlled by the ratio $h_1^u(x, Q^2)/f_1^u(x, Q^2)$ and the scale dependence in this ratio turns out to almost cancel between the numerator and denominator in the valence region relevant at GSI [2]. With GRV and GRSV parameterizations, the ratio $h_1^u(x, 1\text{GeV}^2)/f_1^u(x, 1\text{GeV}^2)$, as well as $g_1^u(x, 1\text{GeV}^2)/f_1^u(x, 1\text{GeV}^2)$, is actually an increasing function of x , leading to the Q -dependence in Fig. 2(b) through $x = Q/\sqrt{s}$. For $\tilde{A}_{LT}$, however, the cancellation of the




FIGURE 2. (a) $\tilde{A}_{LT}$ and (b) $\tilde{A}_{LL}$ and $\tilde{A}_{TT}$ at $x_F = 0$ as a function of Q for $s = 30, 45$ and 210 GeV^2 .

scale dependence between the numerator and denominator in (4) is less complete due to the additional y -integral for the WW part, which, combined with the additional factor x_1 or x_2 ($= Q/\sqrt{s}$), results in the novel Q -dependence in Fig. 2(a).

We have presented a first realistic estimate of the WW contribution to A_{LT} at GSI. Our results indicate that rather large $A_{LT} \sim 10\%$ can be observed. If the strong deviation from our results were observed in the future GSI measurements of A_{LT} , this would provide an indication of large twist-3 effect, associated with the quark-gluon-quark correlation inside nucleon, in particular, in the chiral-odd spin structure. Such data can be analyzed utilizing the QCD evolution of the three-body operators for large number of colors [7], which allows us to avoid the notorious complication in the evolution at higher twist.

ACKNOWLEDGMENTS

The work of Y.K. is supported by Uchida Energy Science Promotion Foundation. The work of K.T. is supported by the Grant-in-Aid for Scientific Research No. B-19340063.

REFERENCES

1. Y. Koike, K. Tanaka and S. Yoshida, Phys. Lett. **B668**, 286 (2008).
2. H. Kawamura, J. Kodaira and K. Tanaka, Phys. Lett. **B662**, 139 (2008).
3. M. Anselmino *et al.*, Phys. Rev. **D75**, 054032 (2007); arXiv:0812.4366 [hep-ph].
4. P.L. Anthony *et al.* [SLAC E155 Collaboration], Phys. Lett. **B553**, 18 (2003).
5. Y. Kanazawa, Y. Koike and N. Nishiyama, Phys. Lett. **B430**, 195 (1998).
6. Y. Koike and K. Tanaka, Phys. Rev. **D51**, 6125 (1995); J. Kodaira, Y. Yasui, K. Tanaka and T. Uematsu, Phys. Lett. **B387**, 855 (1996); J. Kodaira and K. Tanaka, Prog. Theor. Phys. **101**, 191 (1999).
7. A. Ali, V.M. Braun and G. Hiller, Phys. Lett. **B266**, 117 (1991); I.I. Balitsky, V.M. Braun, Y. Koike and K. Tanaka, Phys. Rev. Lett. **77**, 3078 (1996).
8. H. W. Lin, T. Blum, S. Ohta, S. Sasaki and T. Yamazaki, Phys. Rev. **D78**, 014505 (2008).
9. X. Artru and M. Mekhfi, Z. Phys. **C45**, 669 (1990); see also A. Hayashigaki, Y. Kanazawa and Y. Koike, Phys. Rev. **D56**, 7350 (1997); W. Vogelsang, Phys. Rev. **D57**, 1886 (1998).
10. Y. Kanazawa and Y. Koike, Phys. Lett. **B403**, 357 (1997).
11. M. Glück, E. Reya and A. Vogt, Eur. Phys. J. **C5**, 461 (1998).
12. M. Glück, E. Reya, M. Stratmann, and W. Vogelsang, Phys. Rev. **D63**, 094005 (2001).